

**Examination Introduction to Stochastic Processes**  
**(LNMB/Dutch Master Program)**  
**Monday December 19, 2005, 12.00 - 15.00 hours**

This exam consists of 5 problems  
Use of book is not permitted  
Motivate your answers!

1. Suppose that print jobs arrive at a network printer with independent and exponentially distributed inter-arrival times with parameter 10 per hour. It takes the printer exactly 6 seconds to print each page. The sizes of the jobs are i.i.d. and have a Poisson distribution with a mean of 2 pages.
  - a. What is the probability that precisely 20 jobs arrive between 8.30 hours and 10.30 hours?
  - b. What is the probability that a print job arrives while the previous job is not finished, when this previous job consists of 6 pages? (Assume that the printing of this previous job started immediately after its arrival).
  - c. What is the expected arrival time of the first job after 12.00 hours, when the previous job arrived at 11.58 hours?
  - d. Let  $M(t)$  be the number of print jobs consisting of more than 3 pages that arrive in a time interval  $(0, t]$ . Give an expression for  $P(M(t) = m)$ .
  
2. In a simple model, the state of a car dynamo at the beginning of year  $n$  is given by a random variable  $X_n$ , ( $n = 0, 1, 2, \dots$ ), which takes values between 0 and 6. When  $X_n = 0$ , the dynamo is 'as new', when  $X_n = 6$ , it is in 'very bad' condition. Depending on the state  $i$  of the dynamo at the beginning of a year, it breaks down during this year with probability  $i/6$ . After a breakdown, the dynamo is completely revised, so that it is as new at the start of the next year. If a dynamo does not break down during a year, the state is increased by one at the end of the year.
  - a. Show that  $\{X_n, n \geq 0\}$  is a discrete time Markov chain and determine whether the chain is: (i) irreducible, (ii) aperiodic, (iii) transient.
  - b. Give the long run probability that the dynamo is 'as new'.
  - c. Give the long run probability that the dynamo is 'as new', and was in state 2 the year before, i.e. determine  $\lim_{n \rightarrow \infty} P(X_{n-1} = 2, X_n = 0)$ .
  - d. Determine the mean time between two revisions.
  
3. GSM base stations (antenna's) support mobile telephone conversations. Let us consider such a base station and assume that requests for telephone calls arrive according to a Poisson process with rate 30 per hour, while the length of the calls is exponentially distributed with mean 1 minute. Let  $X(t)$  be the number of supported calls at time  $t$ . Then  $\{X(t), t \geq 0\}$  is a continuous-time Markov chain with state space  $\{0, 1, 2, \dots\}$ . Note that when  $X(t) = n$ , all  $n$  calls are in progress simultaneously, so calls are not 'waiting'.

P.T.O.

- a. Let  $n \geq 0$ . Give the distribution of the amount of time that the process spends in state  $n$ , before making a transition into a different state.
- b. Show that the limiting distribution of  $\{X(t), t \geq 0\}$  is Poisson with parameter  $1/2$ .

In reality, base stations can only support a limited number of telephone conversations at the same time. Consider a small base station that can only support 4 conversations (in real stations this capacity is much higher).

- c. From question b., find the long-run probability that the station is completely occupied by using a truncation argument. Which property of  $\{X(t), t \geq 0\}$  do you use?
4. A machine is subject to shocks that occur according to a Poisson process with rate  $\lambda$  per hour. After each shock, the machine breaks down and goes into repair for a stochastic time which is uniformly distributed on  $[0, T]$ . Shocks that occur during repair have no effect. After repair, the machine works properly again. Let  $N(t)$  denote the number of breakdowns in the interval  $(0, t]$ .
    - a. Assuming that a breakdown occurs at time 0, is the process  $\{N(t), t \geq 0\}$  a renewal process? Why (not)?
    - b. On average, how frequently does the machine break down in the long run?
    - c. What is the long run probability that the system is in repair?
  5. Consider a normal renewal process with strictly positive life times  $X_i$ , mean life time  $\mu = EX_1$ , life time LST  $\phi(s) = Ee^{-sX_1}$ , and renewal function  $m(t) = EN(t)$ .
    - a. Let  $S_N(t) = \sum_{i=1}^{N(t)} X_i$  be the time of the last renewal before (or at) time  $t$ , and  $S_{N(t)+1} = \sum_{i=1}^{N(t)+1} X_i$  be the time of the next renewal after time  $t$ . Note that  $X_{N(t)+1} = S_{N(t)+1} - S_{N(t)}$  is the length of the renewal interval containing  $t$ . It is known that  $ES_{N(t)+1} = \mu[m(t) + 1]$ . Can we also argue that  $ES_{N(t)} = \mu m(t)$ ? Explain why or why not.
    - b. The limiting distribution of the excess time is given by  $P(Y \leq x) = \mu^{-1} \int_0^x P(X > y) dy$ . Show that the corresponding Laplace-Stieltjes Transform (LST) is given by  $Ee^{-sY} = (1 - \phi(s))/s\mu$ .

Standard:

1				2				3			4			5		total
a	b	c	d	a	b	c	d	a	b	c	a	b	c	a	b	
1	2	2	3	2	2	2	2	2	3	3	2	2	2	3	3	+ 4 = 40