

# The Sequential Price of Anarchy for Affine Congestion Games with Few Players

Jasper de Jong, Marc Uetz

*University of Twente  
Dept. Applied Mathematics  
7500AE Enschede, The Netherlands*

---

## Abstract

This paper determines the sequential price of anarchy for Rosenthal congestion games with affine cost functions and few players. We show that for two players, the sequential price of anarchy equals 1.5, and for three players it equals  $1039/488 \approx 2.13$ . While the case with two players is analyzed analytically, the tight bound for three players is based on the explicit computation of a worst-case instance using linear programming. The basis for both results are combinatorial arguments to show that finite worst-case instances must exist.

*Keywords:* Congestion Game, Sequential Price of Anarchy  
*2010 MSC:* 91A05, 91A06, 91A46, 90C90

---

## 1. Introduction

Congestion games, introduced by Rosenthal [1], are among the most studied classes of games. In a congestion game, we have  $n$  players  $N = \{1, \dots, n\}$  that compete for certain subsets of a set  $R = \{1, \dots, m\}$  of  $m$  resources, while the  
5 congestion (or cost) that each player experiences per resource, depends on the set of players using it. Subsets of resources that are feasible for a given player  $i \in N$  are given by player specific collections  $\mathcal{A}_i \subseteq 2^R$ . Applications of this class of games include many sorts of allocation problems with scarce resources, and specifically atomic network routing problems, where the resources are network  
10 links and each of the  $n$  players is interested in a least congested path from her origin to destination. Rosenthal congestion games are also referred to as *atomic* congestion games sometimes. This is mainly to distinguish from other types of congestion games such as e.g. the Wardrop traffic models which are also known as non-atomic network routing games, as studied e.g. by Roughgarden  
15 and Tardos [2].

---

*Email address:* `j.dejong-3@utwente.nl`, `m.uetz@utwente.nl` (Jasper de Jong, Marc Uetz)

Rosenthal proved the existence of pure Nash equilibria in any congestion game, by defining what is known as a (Rosenthal) potential function [1]. Assuming that one is interested in some objective function on the set of equilibrium outcomes of a game, the price of anarchy, introduced in 1999 by Koutsoupias and Papadimitriou [3, 4], relates the cost of a worst-case Nash equilibrium to the cost of a globally optimal solution. Shortly after the price of anarchy had been determined for non-atomic network routing games by Roughgarden and Tardos [2], the price of anarchy for atomic congestion games was analyzed as well. Christodoulou and Koutsoupias [5], and independently Awerbuch et al. [6, 7] show that the price of anarchy for congestion games with affine cost functions equals 2 for two players, and  $5/2$  for three or more players.

However, worst-case Nash equilibria can be overly pessimistic in some situations, and specifically in situations where players take their decisions sequentially, worst-case Nash equilibria can often not be realized as equilibrium outcomes. In 2012, Paes Leme et al. [8] have therefore introduced the idea to study the *sequential* price of anarchy. The sequential price of anarchy relates the total cost of the worst-case outcome of a subgame perfect equilibrium of a corresponding extensive form game to the total cost of an optimal solution. This paper establishes tight bounds for the sequential price of anarchy for congestion games with affine cost functions for the case with  $n = 2$  and  $n = 3$  players.

## 2. Related Work and Contribution

Also prior to Paes Leme et al. [8], some authors have addressed the quality of outcomes of network routing games in which players act sequentially. This includes Olver [9], Harks, Heinz and Pfetsch [10], Harks and Vegh [11], as well as Farzad, Olver and Vetta [12]. Without going into details, in these papers the cost of a player depends only on preceding players on the same resource, like in traffic situations where players enter resources in sequence, and a player's cost only depends on the players that entered the resource before. Consequently, players experience different costs on one and the same resource.

We here follow a simpler model where all players experience the same cost per resource. Hence a rational player, if aiming to be farsighted, needs to take into account all future players. This indeed leads to the definition of the sequential price of anarchy as defined by Paes Leme et al. [8]. Subsequent to [8], various researchers studied the sequential price of anarchy for several classes of games, with mixed results. For a handful of problems it was shown that the sequential price of anarchy is strictly smaller than the price of anarchy [8, 13, 14], while for others this turns out to be exactly opposite [15, 16, 17, 18, 19]. Specifically, Correa et al. [19] show that in symmetric, atomic network routing games, the sequential price of anarchy can become as large as  $\Omega(\sqrt{n})$  for large  $n$ ,  $n$  being the number of players, while the price of anarchy is known to be at most  $5/2$  for all  $n \geq 3$  [6, 7]. This of course implies the same lower bound for general congestion games with affine cost functions. Subsequently to [19], work of Groenland and Schäfer [20] suggests that the presence of ties is pivotal for the existence of such worst-case instances. This paper shows that when the number of players is 2

60 or 3, sequential play and subgame perfect equilibria lead to outcomes that are strictly better than the worst-case Nash equilibria. More precisely, we show that the sequential price of anarchy equals 1.5 for two players, and  $1039/488 \approx 2.13$  for three players. Both bounds are shown to be tight. The case with more than three players will be briefly discussed in Section 6.

65 We believe that, although considering the cases of two and three players only, our results are interesting for at least two reasons. First, from an applications point of view it seems not unreasonable to consider (sequential) congestion games with a small number of players, such as the competition for scarce resources by a small number of corporate players, for example. Moreover, through  
70 the lens of computational tractability, sequential games with few players are also the limiting cases that still appear practically feasible. However, note that already for two players, computing the outcome of a subgame perfect equilibrium may be NP-hard [19]. For an arbitrary number of players, it is PSPACE-hard [8]. Second, we obtain our results by using a sequence of combinatorial argu-  
75 ments which yield that the worst-case must be attained on a finite instance of reasonable size. That leads to a linear programming (LP) formulation to actually compute an instance that attains the worst-case for the sequential price of anarchy. In that sense, our proof can be seen as a computerised proof, and the correctness of the result depends on the correctness of the underlying LP; even  
80 though we also verify the resulting worst-case instance. We believe that both the sequence of combinatorial arguments as well as the LP formulation itself are of independent interest, also because it is a first example of doing that for sequential game and subgame perfection.

Inspired by our work, Correa et al. have used another, LP-based proof to  
85 show that for two players, the sequential price of anarchy for atomic *network routing* games is even smaller than that of general two-player congestion games; it equals  $7/5$  [19]. That said, it should be noted that linear programming techniques, and specifically linear programming duality has been used before, e.g. by Nadav and Roughgarden [21] as well as Bilò [22] in order to obtain new or  
90 improved bounds for the price of anarchy in several settings; see also [23].

We end this introductory section by mentioning that -naturally- also more general than affine cost functions have been considered in the context of the price of anarchy for congestion games. A detailed account of all these results is beyond the scope of this paper, and we refer, e.g. to the work by Fotakis [24]  
95 for price of anarchy results with general cost functions.

### 3. Notation and Preliminaries

An instance  $I$  of a *congestion game* is defined as follows: There is a finite set  $R$  of resources, a finite set  $N = \{1, \dots, n\}$  of players, and the action space  $\mathcal{A}_i \subseteq 2^R$  of any player  $i \in N$  is a set of subsets of  $R$ . We say a player  $i$  chooses  
100 a resource  $r \in R$ , if player  $i$  chooses an action  $A_i \ni r$ . Each resource  $r \in R$  has a cost function  $c_r : \mathbb{N} \rightarrow \mathbb{R}^+$  that is nonnegative and nondecreasing. Given an action profile  $A = (A_1, \dots, A_n)$ , each resource  $r$  has cost  $c_r(x_r)$ , where  $x_r$  denotes the number of players choosing  $r$ ,  $x_r = |\{A_i \mid r \in A_i\}|$ . Each player

105  $i \in N$  pays the cost of each of the resources in her chosen action  $A_i$ , which we denote by  $C_i(A) = C_i(A_1, \dots, A_n) = \sum_{r \in A_i} c_r(x_r)$ . The total cost of all players in profile  $A$  is denoted  $C(A) = \sum_{i \in N} C_i(A)$ .

*Affine* congestion games are congestion games where the cost functions of all resources  $r \in R$  are of the form  $c_r(x_r) = \alpha_r + \beta_r x_r$  for  $\alpha_r \geq 0, \beta_r \geq 0$ . Here,  $\alpha_r$  can be thought of as an activation cost for resource  $r$ . We refer to  $\alpha_r$  as the *constant* cost of resource  $r$ , and to  $\beta_r$  as the *weight* of resource  $r$ . We also denote by  $\alpha(R') = \sum_{r \in R'} \alpha_r$  the total constant cost of resources in  $R' \subseteq R$  and by  $\beta(R') = \sum_{r \in R'} \beta_r$  the total weight of resources in  $R' \subseteq R$ .

In a sequential congestion game, let us assume that players act in the order  $1, \dots, n$ . The strategy of player 1 is then one of the available actions from  $\mathcal{A}_1$ . 115 The strategy of a player  $k > 1$ , however, is more complex, as it has to prescribe one action  $A_k \in \mathcal{A}_k$  for each of the possible states that the game can be in, depending on the actions of all players  $i = 1, \dots, k - 1$ . That means that the strategy of player 2 can be written as a vector of actions of length  $|\mathcal{A}_1|$ , for player 3 a vector of actions of length  $|\mathcal{A}_1| \cdot |\mathcal{A}_2|$ , etc. Of course other, more compact representations of players' strategies might be possible, however this is 120 not relevant for what follows. We will use  $S = (S_1, \dots, S_n)$  to denote the profile of strategies of the resulting extensive form game (for a fixed order of players), and  $S^{\text{SPE}}$  to denote a subgame perfect equilibrium in that game.

If we denote by  $A^{\text{OPT}}$  the profile of actions of the  $n$  players in an allocation that minimizes the total cost  $C(A)$ , and by  $A^{\text{SPE}}$  the profile of actions in the outcome of a subgame perfect equilibrium  $S^{\text{SPE}}$ , the *sequential price of anarchy* [8] is defined as

$$\text{SPoA} := \sup \frac{C(A^{\text{SPE}})}{C(A^{\text{OPT}})},$$

125 where the supremum is taken over all possible finite congestion games, all possible permutations of the players, and all possible subgame perfect equilibria.

#### 4. Warmup: The Case of Two Players

We start with the sequential price of anarchy for two players.

**Theorem 1.** *The sequential price of anarchy SPoA equals 1.5 for linear and affine atomic congestion games with two players.*

130 The lower bound is based on the following, simple example.

**Example 2.** *There are two players 1, 2 and three resources 1, 2, 3, with linear cost functions with weights  $\beta_1 = \beta_2 = 1, \beta_3 = 2$ . Player 1 can choose either resource 1 or resource 2. Player 2 can choose either resource 2 or resource 3. This example is illustrated in Figure 1.* ◁

135 For Example 2 it is clearly optimal when player 1 chooses resource 1 and player 2 chooses resource 2, giving a total cost of  $1 + 1 = 2$ . In the outcome of a worst-case subgame perfect equilibrium, however, first player 1 chooses resource 2 and then player 2 chooses resource 3, with a total cost of  $1 + 2 = 3$ . The




Figure 1: The congestion game from Example 2. Dots represent players, squares represent resources, numbers in resources denote weights, and edges represent actions. Fat edges represent chosen actions.

subgame perfect strategy of player 2 is to choose resource 2 if player 1 chooses  
 140 resource 1, and resource 3 if player 1 chooses resource 2. In fact, player 1 is then  
 indifferent, and the worst-case happens if she chooses resource 2. Given that  
 player 1 chooses resource 2, also player 2 is indifferent, and again, the worst-case  
 happens if she chooses resource 3.

Note that this example rests on the existence of ties. However, this can be  
 145 easily avoided, by decreasing the weight of resource 2 by some small constant  
 $\epsilon$  and decreasing the weight of resource 3 by  $3\epsilon$ . Then the above described  
 subgame perfect equilibrium is unique.

We next prove a matching upper bound. The idea is to realize that only a  
 small part of the game tree is relevant, namely the actions that are played in  
 150 some optimal solution, and the actions that are played in equilibrium. That  
 means, we only need to consider *two* actions of player 1, namely optimum and  
 equilibrium, and *three* actions of player 2, namely optimum and the best res-  
 sponses to the two relevant actions of player 1; see also Figure 2.

To formalize this into a proof of the upper bound on the sequential price of  
 155 anarchy, we introduce the following notation. Denote by  $A^{\text{OPT}} = (A_1^{\text{OPT}}, A_2^{\text{OPT}})$   
 the pair of actions of the two players 1 and 2, respectively, in an optimal alloca-  
 tion. Denote by  $A^{\text{SPE}} = (A_1^{\text{SPE}}, A_2^{\text{SPE}})$  the actions in the outcome of a subgame  
 perfect equilibrium. Finally, denote by  $A_2^{\text{SPE}'}$  the subgame perfect action of  
 player 2, in the subgame induced by  $A_1^{\text{OPT}}$  i.e.  $A_2^{\text{SPE}'}$  is a best response of player  
 160 2 to  $A_1^{\text{OPT}}$ . Note that both players might have more actions at their disposal,  
 however these are not relevant for the analysis. Also note that we do not ex-  
 clude cases where any two sets from  $A_1^{\text{OPT}}, A_1^{\text{SPE}}, A_2^{\text{OPT}}, A_2^{\text{SPE}'}$  or  $A_2^{\text{SPE}}$  overlap.  
 It could even be that they are equal. The situation is shown in Figure 2; it  
 displays only the relevant part of the game tree.

Recall that  $\alpha(R')$  and  $\beta(R')$  respectively denote the total constant cost and




Figure 2: All relevant actions in the game tree for 2 player congestion games. Fat lines represent subgame perfect actions.

weight of a set  $R'$  of resources. For brevity, we also use the following notation:

$$\begin{aligned}
 a &= \alpha(A_1^{\text{OPT}}), & b &= \alpha(A_2^{\text{OPT}}), \\
 c &= \beta(A_1^{\text{OPT}} \setminus A_2^{\text{OPT}}), & d &= \beta(A_2^{\text{OPT}} \setminus A_1^{\text{OPT}}), \\
 \gamma &= \beta(A_1^{\text{OPT}} \cap A_2^{\text{OPT}}), & \delta &= \beta(A_1^{\text{OPT}} \cap A_2^{\text{SPE}'}) - \beta(A_1^{\text{OPT}} \cap A_2^{\text{OPT}}).
 \end{aligned}$$

165 Note that  $\delta$  denotes the difference in the total weight of shared resources, when player 1 chooses  $A_1^{\text{OPT}}$  and player 2 switches from  $A_2^{\text{OPT}}$  to  $A_2^{\text{SPE}'}$ .

**Observation 3.**  $C(A^{\text{OPT}}) = C_1(A^{\text{OPT}}) + C_2(A^{\text{OPT}}) = (a + c) + (b + d) + 4\gamma$ .

We prove the upper bound for Theorem 1 by deriving an upper bound on  $C_1(A^{\text{SPE}})$  (Lemma 4) and two upper bounds on  $C_2(A^{\text{SPE}})$  (Lemma 5 and  
170 Lemma 6).

**Lemma 4.**  $C_1(A^{\text{SPE}}) \leq a + c + 2\gamma + \delta$ .

*Proof.*

$$\begin{aligned}
 C_1(A^{\text{SPE}}) &\leq C_1(A_1^{\text{OPT}}, A_2^{\text{SPE}'}) \\
 &= \alpha(A_1^{\text{OPT}}) + \beta(A_1^{\text{OPT}}) + \beta(A_1^{\text{OPT}} \cap A_2^{\text{SPE}'}) \\
 &= a + (c + \gamma) + (\gamma + \delta) \\
 &= a + c + 2\gamma + \delta,
 \end{aligned} \tag{1}$$

where the inequality follows from the Nash inequality (since a subgame perfect equilibrium induces a Nash equilibrium in every state).  $\square$

**Lemma 5.**  $C_2(A^{\text{SPE}}) \leq 2(b + d + \gamma - \delta)$ .

*Proof.*

$$\begin{aligned}
 C_2(A^{\text{SPE}}) &\leq C_2(A_1^{\text{SPE}}, A_2^{\text{SPE}'}) \\
 &\leq \alpha(A_2^{\text{SPE}'}) + 2\beta(A_2^{\text{SPE}'}) \\
 &\leq 2(\alpha(A_2^{\text{SPE}'}) + \beta(A_2^{\text{SPE}'})),
 \end{aligned}$$

where the first inequality follows from the Nash inequality, and the second inequality follows from the fact that each resource can be chosen by at most two players. We now show that  $\alpha(A_2^{\text{SPE}'}) + \beta(A_2^{\text{SPE}'}) \leq b + d + \gamma - \delta$ , proving the lemma. By the Nash inequality, we obtain

$$\begin{aligned} C_2(A_1^{\text{OPT}}, A_2^{\text{SPE}'}) &\leq C_2(A^{\text{OPT}}) \\ &= b + d + 2\gamma \\ &= b + d + \gamma - \delta + (\gamma + \delta). \end{aligned} \quad (2)$$

Since  $\beta(A_1^{\text{OPT}} \cap A_2^{\text{SPE}'}) = \gamma + \delta$ , plugging

$$C_2(A_1^{\text{OPT}}, A_2^{\text{SPE}'}) = \alpha(A_2^{\text{SPE}'}) + \beta(A_2^{\text{SPE}'}) + \beta(A_1^{\text{OPT}} \cap A_2^{\text{SPE}'})$$

into (2), we obtain

$$\alpha(A_2^{\text{SPE}'}) + \beta(A_2^{\text{SPE}'}) \leq b + d + \gamma - \delta. \quad (3)$$

175

□

**Lemma 6.**  $C_2(A^{\text{SPE}}) \leq a + c + b + d + 3\gamma$ .

*Proof.*

$$\begin{aligned} \beta(A_1^{\text{SPE}}) &\leq C_1(A^{\text{SPE}}) \\ &\leq a + c + 2\gamma + \delta, \end{aligned} \quad (4)$$

where the second inequality follows from Lemma 4. Now,

$$\begin{aligned} C_2(A^{\text{SPE}}) &\leq C_2(A_1^{\text{SPE}}, A_2^{\text{SPE}'}) \\ &\leq \alpha(A_2^{\text{SPE}'}) + \beta(A_2^{\text{SPE}'}) + \beta(A_1^{\text{SPE}}) \\ &\leq b + d + \gamma - \delta + a + c + 2\gamma + \delta \\ &= a + c + b + d + 3\gamma. \end{aligned}$$

The first inequality follows from the Nash inequality. The second inequality follows from the fact that each resource that player 1 chooses adds at most the weight of that resource to the cost of player 2. The third inequality follows from (4) and (3). □

180

**Lemma 7.**  $\text{SPoA} \leq 1.5$  for affine atomic congestion games with two players.

*Proof.*

$$\begin{aligned} 2(C(A^{\text{SPE}})) &= 2(C_1(A^{\text{SPE}}) + C_2(A^{\text{SPE}})) \\ &\leq 2a + 2c + 2(2\gamma + \delta) + 2(b + d + \gamma - \delta) + a + c + b + d + 3\gamma \\ &= 3a + 3c + 3b + 3d + 9\gamma \\ &\leq 3C(A^{\text{OPT}}), \end{aligned}$$

where the first inequality follows from Lemmas 4, 5, and 6, and the last inequality follows from Observation 3. □

Theorem 1 now follows from Lemma 7. While the case with two players can  
 185 still be written down algebraically, for three players we turn the same proof idea  
 into a linear programming formulation.

## 5. The Case of Three Players

Along the lines of the proof for the case with two players, we also settle the  
 case with three players.

190 **Theorem 8.**  $\text{SPoA} = \frac{1039}{488} \approx 2.13$  for affine and linear congestion games with  
 three players.

To prove the theorem, we use a linear programming (LP) approach. We  
 first use a sequence of simple, combinatorial arguments to show that a worst-  
 case instance is moderate in size. This is done in Lemmas 9, 10 and 11 below.  
 195 These lemmas apply to games with three players or more. We then compute a  
 worst-case instance for the case with three players using a standard LP solver.

### 5.1. Worst-case instances of moderate size

We use the following notation. Define the series

$$z_1 := 2 \text{ and } z_i := 1 + \prod_{j < i} z_j \text{ for all } i \geq 2.$$

Note that  $z_2 = 3$ ,  $z_3 = 7$ ,  $z_4 = 43$ , and that  $z_i$  grows doubly-exponential.

200 **Lemma 9.** For any instance  $I$  of a congestion game, there exists an instance  
 $I'$ , such that  $|\mathcal{A}_i| \leq z_i$  for all players  $i = 1, \dots, n$ , and  $\text{SPoA}(I') = \text{SPoA}(I)$ .

*Proof.* Denote by  $A^{\text{OPT}}$  an optimal outcome for instance  $I$  and by  $A^{\text{SPE}}$  the  
 outcome of a worst-case subgame perfect equilibrium. The proof goes by suc-  
 cessively eliminating all actions that are not chosen, neither in  $A^{\text{OPT}}$  nor in the  
 subgame perfect equilibrium, in the order of the players  $1, 2, \dots, n$ . For the first  
 205 player, we thereby restrict  $\mathcal{A}_1$  to only two actions:  $A_1^{\text{OPT}}$  and  $A_1^{\text{SPE}}$ . For the sec-  
 ond player, we thereby restrict  $\mathcal{A}_2$  to at most  $z_2 = 3$  actions, namely the actions  
 prescribed by the subgame perfect equilibrium for the two relevant actions of  
 player 1, plus  $A_2^{\text{OPT}}$ . More generally, for the  $k$ -th player, we restrict  $\mathcal{A}_k$  to at  
 most  $1 + \prod_{i < k} z_i$  actions, namely the actions prescribed by the subgame perfect  
 210 equilibrium in each of the at most  $\prod_{i < k} z_i$  states, plus  $A_k^{\text{OPT}}$ . In the so reduced  
 game  $I'$ ,  $S^{\text{SPE}}$  is also subgame perfect, as the actions that were removed are all  
 actions with inferior or identical cost for the respective player.  $\square$

215 **Lemma 10.** For any instance  $I$  of a congestion game, there exists an instance  
 $I'$ , such that  $|\mathcal{A}_i| \leq z_i$  for all players  $i = 1, \dots, n$ , and  $|R| \leq 2^{\sum_{i \in N} |\mathcal{A}_i|} - 1$ , and  
 $\text{SPoA}(I') = \text{SPoA}(I)$ .

*Proof.* By Lemma 9, we may restrict to instances  $I$  with  $|\mathcal{A}_i| \leq z_i$  for all players  $i$ . Suppose the claim is false. Then choose among all instances that falsify the claim an instance  $I$  that minimizes the number of resources  $|R|$ . Then each resource  $r \in R$  is chosen in at least one of the actions. As we have at most  $\sum_{i \in N} |\mathcal{A}_i|$  different actions, there are at most  $2^{\sum_{i \in N} |\mathcal{A}_i|} - 1$  different non-empty subsets of the set of all actions. Since  $|R| > 2^{\sum_{i \in N} |\mathcal{A}_i|} - 1$ , by the pigeonhole principle the “incidence matrix” of all resources and actions contains at least two identical rows. In other words, there must exist two resources  $r, r' \in R$  such that every action either contains both  $r$  and  $r'$ , or it contains neither  $r$  nor  $r'$ . Now we can construct an instance  $I'$  identical to  $I$ , except instead of  $r$  and  $r'$ , it contains a new resource  $r''$  for which  $\beta_{r''} = \beta_r + \beta_{r'}$  and  $\alpha_{r''} = \alpha_r + \alpha_{r'}$ . Other than this replacement, instance  $I'$  has the same sets of actions as instance  $I$ . Now each outcome  $A$  in  $I'$  has the same costs as  $A$  in  $I$ . Therefore the same actions are subgame perfect and  $\text{SPoA}(I') = \text{SPoA}(I)$ . As  $I'$  has one resource less, we obtain a contradiction.  $\square$

Recall that  $A^{\text{OPT}}$  denotes an action profile that minimizes  $C(A)$  for a given instance  $I$ . The final lemma bounds the constant cost and weights in a worst-case instance.

**Lemma 11.** *For any instance  $I$  of a congestion game with affine cost functions, there exists an instance  $I'$  such that  $|\mathcal{A}_i| \leq z_i$  for all players  $i = 1, \dots, n$ ,  $|R| \leq 2^{\sum_{i \in N} |\mathcal{A}_i|} - 1$ ,  $\alpha_r + \beta_r \leq nC(A^{\text{OPT}}) \forall r \in R$ , and  $\text{SPoA}(I') = \text{SPoA}(I)$ .*

*Proof.* We are only left to show the claim on the cost functions. However resources  $r$  with  $\alpha_r + \beta_r > nC(A^{\text{OPT}})$  can safely be eliminated, as it cannot be subgame perfect for any player  $i$  to choose resource  $r$ : choosing  $A_i^{\text{OPT}}$  instead, yields a cost at most  $nC_i(A^{\text{OPT}}) \leq nC(A^{\text{OPT}}) < \alpha_r + \beta_r$ .  $\square$

Specifically, for congestion games with three players, in order to find a worst-case instance we only need to consider games of moderate size. It suffices to let  $|\mathcal{A}_1| = 2, |\mathcal{A}_2| = 3, |\mathcal{A}_3| = 7, |R| = 2^{2+3+7} - 1 = 4095$ , and  $\alpha_r + \beta_r \leq 3C(A^{\text{OPT}})$  for all resources  $r \in R$ . The linear program now works as follows: It maximizes the sequential price of anarchy over all instances with the properties described above. We have  $2^{12} - 1 = 4095$  resources, one for every (potential) nonempty intersection of actions. The LP decides the weight  $\beta_r \geq 0$  and constant cost  $\alpha_r \geq 0$  of each resource. We define some fixed outcome as the optimum solution with total cost normalized to 1, which yields that  $\alpha_r + \beta_r \leq 3$  for all  $r \in R$ , and we maximize the sequential price of anarchy.

## 5.2. Details LP formulation

In violation to our earlier nomenclature, we here choose to denote actions by lowercase letters  $a, a', b, b', c, c', \mu$  and  $\nu$ , because they appear as index to the decision variables. We use binary parameters  $\delta_{\mu r}$  to specify whether resource  $r$  is chosen in action  $\mu$ . For each resource  $r$ , we have decision variables  $\alpha_r$  and  $\beta_r$ , the constant cost and weight of  $r$ , respectively. We fix a subgame perfect equilibrium  $S^{\text{SPE}}$  with subgame perfect outcome  $A^{\text{SPE}}$  and we use binary

parameters  $z_a^1, z_{ab}^2$  and  $z_{abc}^3$  to prescribe which actions are part of the subgame perfect equilibrium  $S^{\text{SPE}}$  for players 1, 2, 3 respectively. To clarify these, for example  $z_{ab}^2 = 1$  means that action  $b$  is the subgame perfect action by player 2, given player 1 plays  $a$ . We denote by  $i.x$  the  $x$ -th action of player  $i$ . We define the action profile  $(1.1, 2.1, 3.1)$  as the optimal outcome with total cost normalized to 1. The subgame perfect equilibrium is also fixed, namely for player 1 it is action 1.2, for player 2 it is action 2.2 (given 1.1) and action 2.3 (given 1.2). For player 3, in branch  $a \geq 1$  of the game tree, we let action  $3.(a+1)$  be the subgame perfect action. This implies that action profile  $(1.2, 2.3, 3.7)$  is the outcome of the worst-case subgame perfect equilibrium.

We denote by  $v_\mu = \sum_{r \in R} \delta_{\mu r} (\beta_r + \alpha_r)$  the cost of a player who chooses action  $\mu$ , assuming no other players would be there. Next, we denote by  $o_{\mu\nu} = \sum_{r \in R} \delta_{\mu r} \delta_{\nu r} \beta_r$  the additional costs that two players with actions  $\mu$  and  $\nu$  incur due to overlap in resources. We will use these auxiliary variables to determine the total cost of player  $i$  when players 1, 2 and 3 choose actions  $a, b$  and  $c$ , respectively. This we denote by  $C_i(abc)$ .

Since we have to be able to describe a subgame perfect equilibrium, we need a bit more of notation:  $C_1(a)$  and  $C_2(ab)$  denote the cost of actions of players 1 and 2 respectively, when successors play subgame perfect. For instance,  $C_2(ab)$  denotes the cost of action  $b$  for player 2, when player 1 chooses action  $a$  and player 3 plays subgame perfect. Finally  $C(A^{\text{SPE}})$  will denote the total costs for all players in the subgame perfect outcome.

For each of the following parameters, variables, and constraints, we assume the following conventions on nomenclature:

$$a, a' \in \mathcal{A}_1, b, b' \in \mathcal{A}_2, c, c' \in \mathcal{A}_3, \mu, \nu \in \cup_{i \in N} \mathcal{A}_i, r \in R, i \in N.$$

In writing,  $a$  and  $a'$  denote actions of player 1,  $b$  and  $b'$  denote actions of player 2,  $c$  and  $c'$  denote actions of player 3,  $\mu$  and  $\nu$  are arbitrary actions,  $r$  is a resource, and  $i$  is a player. Also, let us denote by  $S^{\text{SPE}}$  the worst-case subgame perfect equilibrium that we seek to compute, so that  $S_i^{\text{SPE}}$  is the strategy of player  $i = 1, 2, 3$ . Also recall that  $A^{\text{SPE}} = (1.2, 2.3, 3.7)$  is the resulting action profile. The linear program is now as follows.

*Binary parameters*

$$\begin{array}{ll} \delta_{\mu r} & \forall r, \mu \begin{cases} 1 & \text{if } r \in \mu \\ 0 & \text{otherwise} \end{cases} \\ z_a^1 & \forall a \begin{cases} 1 & \text{if } a \text{ is prescribed by } S_1^{\text{SPE}} \\ 0 & \text{otherwise} \end{cases} \\ z_{ab}^2 & \forall a, b \begin{cases} 1 & \text{if } b \text{ is prescribed by } S_2^{\text{SPE}} \text{ in state } a \\ 0 & \text{otherwise} \end{cases} \\ z_{abc}^3 & \forall a, b, c \begin{cases} 1 & \text{if } c \text{ is prescribed by } S_3^{\text{SPE}} \text{ in state } ab \\ 0 & \text{otherwise} \end{cases} \end{array}$$

*Variables*

$\alpha_r$	$\forall r$	constant cost of $r$
$\beta_r$	$\forall r$	weight of $r$
$v_\mu$	$\forall \mu$	constant cost plus weight of resources in $\mu$
$o_{\mu\nu}$	$\forall \mu, \nu   \mu \neq \nu$	weight of resources in $\mu \cap \nu$
$C_i(abc)$	$\forall a, b, c, i$	cost of player $i$ when players 1,2,3 choose $a, b, c$ respectively
$C(A^{\text{SPE}})$		costs in subgame perfect outcome $A^{\text{SPE}}$
$C_1(a)$	$\forall a$	cost of player 1 when she chooses $a$ and 2, 3 choose as prescribed by $S^{\text{SPE}}$
$C_2(ab)$	$\forall a, b$	costs of player 2 when players 1, 2 choose $a, b$ and 3 chooses as prescribed by $S^{\text{SPE}}$

290 *Constraints*

$$0 \leq \alpha_r + \beta_r \leq 3 \quad \forall r \quad (5)$$

$$v_\mu = \sum_{r \in R} \delta_{\mu r} (\beta_r + \alpha_r) \quad \forall \mu \quad (6)$$

$$o_{\mu\nu} = \sum_{r \in R} \delta_{\mu r} \delta_{\nu r} \beta_r \quad \forall \mu, \forall \nu | \mu \neq \nu \quad (7)$$

$$C_1(abc) = v_a + o_{ab} + o_{ac} \quad \forall a, b, c \quad (8)$$

$$C_2(abc) = v_b + o_{ab} + o_{bc} \quad \forall a, b, c \quad (9)$$

$$C_3(abc) = v_c + o_{ac} + o_{bc} \quad \forall a, b, c \quad (10)$$

$$C_3(abc) \leq C_3(abc') \quad \forall a, b, c | z_{abc}^3 = 1, c' \quad (11)$$

$$C_2(ab) \leq C_2(ab') \quad \forall a, b | z_{ab}^2 = 1, b' \quad (12)$$

$$C_1(a) \leq C_1(a') \quad \forall a | z_a^1 = 1, a' \quad (13)$$

$$C_1(a) = C_1(abc) \quad \forall a, b | z_{ab}^2 = 1, c | z_{abc}^3 = 1 \quad (14)$$

$$C_2(ab) = C_2(abc) \quad \forall a, b, c | z_{abc}^3 = 1 \quad (15)$$

$$\sum_{i \in N} C_i(1.1, 2.1, 3.1) = 1 \quad (16)$$

$$C(A^{\text{SPE}}) = \sum_{i \in N} C_i(1.2, 2.3, 3.7) \quad (17)$$

Constraints (6) and (7) define costs of actions  $v_\mu$  and intersection costs  $o_{\mu\nu}$  for all actions  $\mu, \nu$ . Constraints (8),(9), and (10) define the costs in each outcome for each player. Constraints (11),(12), and (13) guarantee that no player can




Figure 3: Example 12 for three players. Squares represent resources. The number in each resource denotes its weight. Encircled resources depict the actions.

improve from the subgame perfect equilibrium  $S^{\text{SPE}}$ . Constraints (14) define  
 295  $C_1(a)$ , and constraints (15) define  $C_2(ab)$ , as explained earlier. The optimal  
 solution is the action profile (1.1, 2.1, 3.1) and has total cost equal to 1, encoded  
 into constraint (16). Constraint (17) defines  $C(A^{\text{SPE}})$ . Finally, constraints (5)  
 are not necessary, but bound the coefficients  $\alpha_r$  and  $\beta_r$ .

The objective is to maximize  $C(A^{\text{SPE}})$ , since, due to the normalization, this  
 300 value equals the sequential price of anarchy. This allows us to give the LP-based  
 proof of the main theorem of this paper. This result is particularly interesting  
 in comparison to the tight bound 2.5 for the price of anarchy for non-sequential  
 three player congestion games [6, 7, 5].

*Proof of Theorem 8.* We have solved the above described linear program using  
 305 the AIMMS modeling framework with CPLEX 12.5 as LP solver. We obtain an  
 optimal solution with value  $\frac{1039}{488} \approx 2.13$ , which proves Theorem 8.  $\square$

### 5.3. Tight Lower Bound Instance

We have used a mixed integer linear program (MIP) in order to model situations with smaller action spaces and fewer resources. Here, we use integer variables (instead of parameters) to decide which actions are subgame perfect for the third player. Naturally, such a MIP only yields lower bounds on the sequential price of anarchy, but no upper bounds. Nevertheless, after inspection of the solution of the linear program, using the MIP we have been able to compute a lower bound example that matches the upper bound of Theorem 8, so that player 3 only uses four different actions. We here give this instance, scaled to integers such that the cost in the optimum solution equals 488.

**Example 12.** *There are three players. Player 1 has two actions 1.1, 1.2. Player 2 has three actions 2.1, 2.2, 2.3. Player 3 has 4 actions 3.1, 3.2, 3.3, 3.4. There are 13 resources  $R = \{1, \dots, 13\}$  with constant costs  $\alpha_r = 0$  for all  $r \in R$ . Table 1 shows for each resource, its weight and the actions that contain it. This instance is also illustrated in a more intuitive way in Figure 3.*  $\triangleleft$

$r$	$\beta_r$	1.1	1.2	2.1	2.2	2.3	3.1	3.2	3.3	3.4
1	84		✓		✓		✓			
2	52		✓		✓	✓	✓			
3	3		✓			✓	✓		✓	
4	4		✓		✓	✓	✓	✓		
5	31				✓	✓	✓	✓		
6	52		✓	✓					✓	
7	54			✓		✓		✓		
8	92	✓				✓		✓	✓	
9	51	✓				✓			✓	
10	28		✓	✓						
11	4		✓	✓				✓	✓	
12	33			✓				✓		✓
13	374									✓

Table 1: The weights and actions of each resource. ✓ denotes that the corresponding action contains the corresponding resource.

Backward induction on the game tree yields a subgame perfect outcome where player 1 chooses 1.2, player 2 chooses 2.3, and player 3 chooses 3.4, yielding a total cost of  $C(A^{\text{SPE}}) = 1039$ . The game tree is shown in Figure 4.

## 6. Conclusions

The linear program that we proposed here for the case of three players, is no longer practically feasible for the case with four players, as the number of actions of the fourth player would be 43, and the number of resources would become  $2^{55} - 1$ . Yet we have experimented with the reduced size MIP in order to compute




Figure 4: The game tree of Example 12 for three players. The number at each action denotes the cost of the corresponding player when all successors play subgame perfect. Fat lines are actions of the subgame perfect equilibrium.

330 lower bounds on the sequential price of anarchy for congestion games with four  
 or more players. Even though the corresponding MIP was large and posed some  
 computational challenges, we have been able to show that  $\text{SPoA} > 2.55$  for four  
 players [25], so specifically it is larger than  $5/2$  (which is the price of anarchy  
 for Nash equilibria).

### 335 Acknowledgements

We thank José Correa and Bart de Keijzer for helpful discussions. An ex-  
 tended abstract of this paper has been published in the proceedings of the 10th  
 Conference on Web and Internet Economics [26]. Jasper de Jong was supported  
 by 3TU.AMI.

### 340 References

- [1] R. W. Rosenthal, A class of games possessing pure-strategy Nash equilibria, *Internat. J. Game Theory* 2 (1) (1973) 65–67.
- [2] T. Roughgarden, É. Tardos, How bad is selfish routing?, *Journal of the ACM* 49 (2) (2002) 236–259.
- 345 [3] E. Koutsoupias, C. Papadimitriou, Worst-case equilibria, in: C. Meinel, S. Tison (Eds.), 16th Annual Symposium on Theoretical Aspects of Computer Science (STACS 1999), Vol. 1563 of Lecture Notes in Computer Science, Springer, 1999, pp. 404–413.
- [4] E. Koutsoupias, C. Papadimitriou, Worst-case equilibria, *Computer Science Re-*  
 350 *view* 3 (2009) 65–69.
- [5] G. Christodoulou, E. Koutsoupias, The price of anarchy of finite congestion games, in: *Proc. 37th Annual ACM Symp. on the Theory of Computing (STOC 2005)*, 2005, pp. 67–73.
- 355 [6] B. Awerbuch, Y. Azar, A. Epstein, The price of anarchy of routing unsplittable flow, in: *Proc. 37th Annual ACM Symp. on the Theory of Computing (STOC 2005)*, 2005, pp. 57–66.

- [7] B. Awerbuch, Y. Azar, A. Epstein, The price of routing unsplittable flow, *SIAM Journal on Computing* 42 (1) (2013) 160–177.
- [8] R. Paes Leme, V. Syrgkanis, É. Tardos, The curse of simultaneity, in: Proc. 3rd Conf. on Innovations in Theoretical Computer Science (ICTS 2012), ACM, 2012, pp. 60–67.
- [9] N. Olver, The price of anarchy and a priority-based model of routing, Master’s thesis, McGill University, Montréal, Canada (2006).
- [10] T. Harks, S. Heinz, M. Pfetsch, Competitive online multicommodity routing, *Theory of Computing Systems* 45 (2009) 533–545.
- [11] T. Harks, L. Vegh, Selfish routing with online demands, in: J. Jansen, P. Pralat (Eds.), Proc. 4th Workshop on Combinatorial and Algorithmic Aspects of Networking (CAAN 2007), Vol. 4852 of Lecture Notes in Computer Science, Springer, 2007, pp. 27–45.
- [12] B. Farzad, N. Olver, A. Vetta, A priority-based model of routing, *Chicago Journal of Theoretical Computer Science* 2008 (1).
- [13] J. de Jong, M. Uetz, A. Wombacher, Decentralized throughput scheduling, in: P. Spirakis, M. Serna (Eds.), Algorithms and Complexity (CIAC 2013), Vol. 7878 of Lecture Notes in Computer Science, Springer, 2013, pp. 134–145.
- [14] R. Hassin, U. Yovel, Sequential scheduling on identical machines, *Operations Research Letters* 43 (5) (2015) 530–533.
- [15] V. Bilò, M. Flammini, G. Monaco, L. Moscardelli, Some anomalies of farsighted strategic behavior, in: C. Kaklamakis, K. Pruhs (Eds.), Approximation and Online Algorithms (WAOA 2010), Vol. 8447 of Lecture Notes in Computer Science, Springer, 2013, pp. 229–241.
- [16] V. Bilò, M. Flammini, G. Monaco, L. Moscardelli, Some anomalies of farsighted strategic behavior, *Theory of Computing Systems* 56 (1) (2015) 156–180.
- [17] A. Angelucci, V. Bilò, M. Flammini, L. Moscardelli, On the sequential price of anarchy of isolation games, in: D.-Z. Du, G. Zhang (Eds.), Computing and Combinatorics (COCOON 2013), Vol. 7936 of Lecture Notes in Computer Science, Springer, 2013, pp. 17–28.
- [18] A. Angelucci, V. Bilò, M. Flammini, L. Moscardelli, On the sequential price of anarchy of isolation games, *Journal of Combinatorial Optimization* 29 (1) (2015) 165–181.
- [19] J. R. Correa, J. de Jong, B. de Keijzer, M. Uetz, The curse of sequentiality in routing games, in: E. Markakis, G. Schäfer (Eds.), Web and Internet Economics (WINE 2015), Vol. 9470 of Lecture Notes in Computer Science, Springer, 2015, pp. 258–271.
- [20] C. Groenland, G. Schäfer, The curse of ties in congestion games with limited lookahead, in: Proceedings of the 17th International Conference on Autonomous Agents and Multiagent Systems (AAMAS 2018), IFAAMAS, 2018, pp. 1941–1943.

- 400 [21] U. Nadav, T. Roughgarden, The limits of smoothness: a primal-dual framework for price of anarchy bounds, in: A. Saberi (Ed.), *Internet and Network Economics (WINE 2010)*, Vol. 6484 of *Lecture Notes in Computer Science*, Springer, 2010, pp. 319–326.
- [22] V. Bilò, A unifying tool for bounding the quality of non-cooperative solutions in weighted congestion games, *Theory of Computing Systems* 62 (5) (2018) 1288–1317.
- 405 [23] V. Bilò, C. Vinci, On stackelberg strategies in affine congestion games, in: E. Markakis, G. Schäfer (Eds.), *Web and Internet Economics (WINE 2015)*, Vol. 9470 of *Lecture Notes in Computer Science*, Springer, 2015, pp. 132–145.
- [24] D. Fotakis, Stackelberg strategies for atomic congestion games, *Theory of Computing Systems* 47 (2010) 218–249.
- 410 [25] K. Kolev, Sequential price of anarchy for atomic congestion games with limited number of players, Master’s thesis, University of Twente (2016).
- [26] J. de Jong, M. Uetz, The sequential price of anarchy for atomic congestion games, in: T.-Y. Liu, Q. Qi, Y. Ye (Eds.), *Web and Internet Economics (WINE 2014)*, Vol. 8877 of *Lecture Notes in Computer Science*, Springer, 2014, pp. 429–434.